



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

elder seeds. Their habits of feeding were exactly as given by Mr. Keyes in the January 'Auk' (p. 114). The remaining flock of six females stayed for three weeks and then departed.—GEO. A. COLEMAN, *London, Nemaha Co., Nebraska.*

**Tameness of the Pine Siskin.**—On April 29, 1888, while walking near Oak Hill in Newton, Mass., I noticed two Pine Siskins (*Spinus pinus*), about a heap of hops by the roadside. One of them flew away at my approach, but the other remained there feeding, and, though perfectly able-bodied and in good condition, was remarkably tame. I stood watching him some time. After a while I reached out and stroked him, and finally succeeded in catching him in one hand. When I let him go, he flew off to some distance. Before I caught him, he went and perched in a bush near by and apparently went to sleep, putting his head over his left wing under the scapular feathers, so that it was completely hidden. When I approached too near, he would take his head out and look at me and then put it back again when I drew back. The ground about there was sprinkled with droppings, showing that the birds had probably been there for some time. Was this bird affected by the hops, or is there any other explanation of his curious conduct? The hops were to be used as dressing for a field of grass. —FRANCIS H. ALLEN, *West Roxbury, Mass.*

**Further Notes on Seaside Sparrows.**—A series of six specimens of Seaside Sparrows kindly loaned me for examination by Mr. G. S. Miller, Jr., of Peterboro, N. Y., includes four specimens from Sapelo Island, coast of Georgia, one from Cedar Keys, and one from Corpus Christi, Texas. The Corpus Christi specimen (male, May 26, 1886) is typically *Ammodramus maritimus sennetti*; the Cedar Keys example (female, Jan. 30, 1880) is typical *A. m. peninsulae*, as is also one of the four specimens from Sapelo Island, the other three being *A. maritimus*. All of the Sapelo Island specimens were taken in December, examples of both forms being labelled Dec. 14, 1887. Mr. Miller kindly wrote me concerning the Sapelo specimens before sending them, as follows: "They all seem to be true *A. maritimus*, excepting one female taken Dec. 14, which is, so far as I can see, typical *peninsulae*. It agrees in almost every particular with a specimen taken at Cedar Keys, Fla., which I should refer without hesitation to this form. Should the Sapelo Island specimen prove to be *peninsulae*, it would extend the range of that form considerably."—J. A. ALLEN, *Am. Mus. Nat. Hist., New York City.*

**A Second Instance of the Breeding of the White-throated Sparrow in Eastern Massachusetts.**—On the 13th and 14th of June (1888) I found a White-throated Sparrow (*Zonotrichia albicollis*) singing in Wakefield, Mass.; and on the 16th and 17th, in the same spot (a bushy roadside swamp), I saw a female of the same species. The male was in full plumage, and the identification was absolute in every case. I saw nothing more of either bird, as I left home on the 18th and did not return